

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/SE05/000501

International filing date: 06 April 2005 (06.04.2005)

Document type: Certified copy of priority document

Document details: Country/Office: SE
Number: 0400969-2
Filing date: 14 April 2004 (14.04.2004)

Date of receipt at the International Bureau: 26 April 2005 (26.04.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

PRV

PATENT- OCH REGISTRERINGSVERKET
Patentavdelningen

PCT / SE 2005 / 0 0 0 5 0 1

Intyg Certificate

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

Ansökan ingavs ursprungligen på engelska.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

The application was originally filed in English.


(71) Sökande AstraZeneca AB, Södertälje SE
Applicant (s)

(21) Patentansökningsnummer 0400969-2
Patent application number

(86) Ingivningsdatum 2004-04-14
Date of filing

Stockholm, 2005-04-19

För Patent- och registreringsverket
For the Patent- and Registration Office


Görel Gustafsson

Avgift
Fee

PATENT- OCH
REGISTRERINGSVERKET
SWEDEN

Postadress/Adress
Box 5055
S-102 42 STOCKHOLM

Telefon/Phone
+46 8 782 25 00
Vx 08-782 25 00

Telex
17978
PATOREG S

Telefax
+46 8 666 02 86
08-666 02 86

ARYL GLYCINE DERIVATIVES AND THEIR USE**FIELD OF THE INVENTION**

This invention relates to the treatment of diseases in which serotonin, Substance P or Neurokinin A are implicated, for example, in the treatment of disorders or conditions such as hypertension, depression, generalized anxiety disorder, phobias, posttraumatic stress syndrome, avoidant personality disorder, premature ejaculation, eating disorders, obesity, chemical dependencies, cluster headache, migraine, pain, Alzheimer's disease, obsessive-compulsive disorder, panic disorder, memory disorders, Parkinson's disease, endocrine disorders vasospasm, cerebellar ataxia, gastrointestinal tract disorders, negative symptoms of schizophrenia, premenstrual syndrome, fibromyalgia syndrome, stress incontinence, Tourette's syndrome, trichotillomania, kleptomania, male impotence, attention deficit hyperactivity disorder, chronic paroxysmal hemicrania and headache.

BACKGROUND

The mammalian neurokinins are peptide neurotransmitters found in the peripheral and central nervous systems. The three principal neurokinins are Substance P (SP), Neurokinin A (NKA) and Neurokinin B (NKB). N-terminally extended forms of at least NKA are known. Three receptor types are known for the principal neurokinins. Based upon their relative selectivities for the neurokinins SP, NKA and NKB, the receptors are classified as neurokinin 1 (NK₁), neurokinin 2 (NK₂) and neurokinin 3 (NK₃) receptors, respectively. In the periphery, SP and NKA are localized in C-afferent sensory neurons, which neurons are characterized by non-myelinated nerve endings known as C-fibers, and are released by selective depolarization of these neurons, or selective stimulation of the C-fibers. C-Fibers are located in the airway epithelium, and the tachykinins are known to cause profound effects which clearly parallel many of the symptoms observed in asthmatics. The effects of release or introduction of tachykinins in mammalian airways include bronchoconstriction, increased microvascular permeability, vasodilation, increased mucus secretion and activation of mast cells. Neurokinin antagonists that interact with NK₁, NK₂ and NK₃ receptors, having different chemical structures have been described. Particularly international publications WO 98/07722, WO 96/39383 and WO 98/25617, and regional publications EP 428434, EP 474561, EP 515240 and EP 559538 disclose the preparation of a variety of chemical structures.


NK₁ activity is also implicated in depression and anxiety, mice with genetically altered NK₁ receptors have decreased anxiety related behavior (Santarelli, L., *et. al.*, Proc. Nat. Acad. Sci. (2001), 98, 1912) and NK₁ antagonists have been reported to be effective in an animal model of depression (Papp, M., *et. al.*, Behav. Brain Res. (2000), 115, 19).

5 Selective Serotonin Reuptake Inhibitors (SSRIs) are widely used for the treatment of major depressive disorder (MDD) and are considered well-tolerated and easily administered. SSRIs, however, have a delayed onset of action, are associated with undesirable side effects, such sexual dysfunction, and are ineffective in perhaps 30% of patients (M. J. Gitlin, MJ, J. Clin. Psych., 55, 406-413, 1994).

10 Compounds with dual action as NK₁ antagonists and serotonin reuptake inhibitors may also provide a new class of antidepressants. Indeed, compounds combining NK₁ antagonism and serotonin reuptake inhibition have been described (Ryckmans, T., *et. al.*, Bioorg. Med. Chem. Lett. (2002), 12, 261).

SUMMARY OF THE INVENTION

15 The present invention encompasses compounds having neurokinin 1 ("NK₁") antagonist activity and/or serotonin reuptake inhibitory ("SRI") activity. Aryl glycine compounds of the invention are those on accord with formula I



I

wherein:

20 R¹ and R² are independently selected from C₁₋₆ alkyl or C₁₋₆ alkenyl, or together with the N to which they are bound, form heterocycle containing 3-6 atoms or heterocycle substituted with moieties independently selected from hydrogen, halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy or C₁₋₄ alkyl substituted with 1, 2 or 3 halo moieties, amino, or amino substituted with C₁₋₄ alkyl, C₁₋₄ alkoxy or C₁₋₄ alkyl, substituted with 0, 1, 2, or 3 halo moieties; in vivo-hydrolysable precursors thereof, and pharmaceutically-acceptable salts thereof;

R³ is selected from hydrogen or C₁₋₆ alkyl;

X is selected from O or NR⁴;

R⁴ is selected from hydrogen or C₁₋₆ alkyl;

Y is C₁₋₆ alkyl or C(O);

n is 0, 1 or 2;

- 5 Ar¹ is phenyl or phenyl substituted with moieties independently selected from hydrogen, halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy or C₁₋₄ alkyl substituted with 1, 2 or 3 halo moieties; in vivo-hydrolysable precursors thereof, and pharmaceutically-acceptable salts thereof; and

- Ar² phenyl, naphthyl, tetralin, or phenyl, naphthyl or tetralin substituted with moieties independently selected from hydrogen, halogen, cyano, nitro, C₁₋₄ alkyl, C₁₋₄ alkoxy or C₁₋₄ alkyl substituted with 1, 2 or 3 halo moieties; in vivo-hydrolysable precursors thereof, and pharmaceutically-acceptable salts thereof.


- The invention also encompasses in vivo-hydrolysable precursors and pharmaceutically-acceptable salts of the compounds, pharmaceutical compositions and formulations containing them, methods of using them to treat diseases and conditions either alone or in combination with other therapeutically-active compounds or substances, processes and intermediates used to prepare them, uses of them as medicaments, uses of them in the manufacture of medicaments and uses of them for diagnostic and analytic purposes.

DETAILED DESCRIPTION OF THE INVENTION

Compounds of this invention have NK₁ antagonist activity and/or SRI activity.

- 20 Accordingly, this invention comprises such compounds, pharmaceutical compositions containing such compounds and methods of using such compounds to treat central nervous system (CNS) and other disorders.

Compounds of the invention are those in accord with formula I:



I

wherein:

R^1 and R^2 are independently selected from C_{1-6} alkyl or C_{1-6} alkenyl, or together with the N to which they are bound, form heterocycle containing 3-6 atoms or heterocycle substituted with moieties independently selected from hydrogen, halogen, C_{1-4} alkyl, C_{1-4} alkoxy or C_{1-4} alkyl substituted with 1, 2 or 3 halo moieties, amino, or amino substituted with C_{1-4} alkyl, C_{1-4} alkoxy or C_{1-4} alkyl, substituted with 0, 1, 2, or 3 halo moieties; in vivo-hydrolysable precursors thereof, and pharmaceutically-acceptable salts thereof;

R^3 is independently selected from hydrogen or C_{1-6} alkyl;

X is selected from O or NR^4 ;

R^4 is selected from hydrogen or C_{1-6} alkyl;

Y is C_{1-6} alkyl or $C(O)$;

n is 0, 1 or 2;

Ar^1 is phenyl or phenyl substituted with moieties independently selected from hydrogen, halogen, C_{1-4} alkyl, C_{1-4} alkoxy or C_{1-4} alkyl substituted with 1, 2 or 3 halo moieties; in vivo-hydrolysable precursors thereof, and pharmaceutically-acceptable salts thereof; and

Ar^2 phenyl, naphthyl, tetralin, or phenyl, naphthyl or tetralin substituted with moieties independently selected from hydrogen, halogen, cyano, nitro, C_{1-4} alkyl, C_{1-4} alkoxy or C_{1-4} alkyl substituted with 1, 2 or 3 halo moieties; in vivo-hydrolysable precursors thereof, and pharmaceutically-acceptable salts thereof.

Particular compounds of the invention are those of the examples herein.

Another aspect of the invention is pharmaceutically-acceptable salts of a compounds as described herein made with an inorganic or organic acid which affords a physiologically-acceptable anion.

Particular pharmaceutically-acceptable salts of compounds of the invention are those wherein the inorganic or organic acid is selected from hydrochloric, hydrobromic, sulfuric, phosphoric, methanesulfonic, sulfamic, para-toluenesulfonic, acetic, citric, lactic, tartaric, malonic, fumaric, ethanesulfonic, benzenesulfonic, cyclohexylsulfamic, salicylic and quinic acids.

Another aspect of the invention is a pharmaceutical composition comprising a compound of the invention or an in vivo-hydrolysable precursor or a pharmaceutically-acceptable salt thereof and a pharmaceutically-acceptable carrier.

Yet another aspect of the invention is a method of treating a disease condition wherein antagonism of NK₁ receptors and/or SRI activity is beneficial which method comprises administering to a warm-blooded animal an effective amount of a compound of the invention or an in vivo-hydrolysable precursor or a pharmaceutically-acceptable salt thereof.

5 Still another aspect of the invention is the use of a compound of the invention or an in vivo-hydrolysable precursor or a pharmaceutically-acceptable salt thereof in the preparation of a medicament for use in a disease condition wherein antagonism of the NK₁ receptors and/or SRI activity is beneficial.

A further aspect of the invention is a method for treating a disorder or condition selected
10 from hypertension, depression in cancer patients, depression in Parkinson's patients, postmyocardial infarction depression, subsyndromal symptomatic depression, depression in infertile women, pediatric depression, major depression, single episode depression, recurrent depression, child abuse induced depression, post partum depression, generalized anxiety disorder, agoraphobia, social phobia, simple phobias, posttraumatic stress syndrome, avoidant personality
15 disorder, premature ejaculation, anorexia nervosa, bulimia nervosa, obesity, addictions to alcohol, cocaine, heroin, phenobarbital, nicotine or benzodiazepines; cluster headache, migraine, pain, Alzheimer's disease, obsessive-compulsive disorder, panic disorder, dementia, amnestic disorders, age-related cognitive decline, dementia in Parkinson's disease, neuroleptic-induced parkinsonism, tardive dyskinesias, hyperprolactinaemia, vasospasm, cerebral vasculature
20 vasospasm, cerebellar ataxia, gastrointestinal tract disorders, negative symptoms of schizophrenia, premenstrual syndrome, fibromyalgia syndrome, stress incontinence, Tourette's syndrome, trichotillomania, kleptomania, male impotence, attention deficit hyperactivity disorder, chronic paroxysmal hemicrania and headache associated with vascular disorders in a mammal, wherein antagonism of the NK₁ receptors and SRI activity is beneficial, comprising
25 administering an effective amount of a compound of the invention or a pharmaceutically-acceptable salt thereof effective in treating such disorder or condition.

In a particular aspect of the invention the method for treating a disorder or condition mentioned herein, comprises administering a compound of the invention in combination with a pharmaceutically-acceptable carrier.

30 Compounds in accord with formula I and their in vivo-hydrolysable precursors or a pharmaceutically-acceptable salts may be made by processes as described and exemplified herein

and by processes similar thereto and by processes known in the chemical art. If not commercially available, starting materials for these processes may be made by procedures which are selected from the chemical art using techniques which are similar or analogous to the synthesis of known compounds.

5 Pharmaceutically-acceptable salts may be prepared from the corresponding acid in a conventional manner. Non-pharmaceutically-acceptable salts may be useful as intermediates and as such are another aspect of the present invention.

It is well known in the art how to prepare optically-active forms (for example, by resolution of the racemic form or by synthesis from optically-active starting materials) and all
10 optically active forms, enantiomers are compounds of this invention. Further, the mixture of enantiomers can have either or both serotonin and neurokinin activity and/or either of the pure enantiomers can have serotonin and or neurokinin activity.

The following biological test methods, data and Examples serve to illustrate and further describe the invention.

15 The utility of a compound of the invention or an in vivo-hydrolysable precursor or a pharmaceutically-acceptable salt thereof (hereinafter, collectively referred to as a "Compound") may be demonstrated by standard tests and clinical studies, including those disclosed in the publications described below.

Biological Assays:

20 Test A — SERT Binding Assay:

Frozen membrane preparations of a stably transfected HEK293 cell line expressing human 5-HTT receptors were purchased from Receptor Biology (PerkinElmer). Frozen aliquots were rapidly thawed, homogenized, and diluted in assay buffer (AB) containing 50 mM TRIS-HCL, 120 mM NaCl, 5 mM KCl and adjusted to pH 7.4 with NaOH. Final protein concentration
25 was 40 µg/ml. Test compounds were evaluated in competition assays utilizing [³H]-Imipramine Hydrochloride purchased from NEN (PerkinElmer) as the radioligand. The stock radioligand was diluted with AB for a final concentration of approximately 2 nM. K_d for [³H]-Imipramine Hydrochloride was determined to be 2.7 nM. The competition assays were performed on 96-well assay plates — two drugs per plate. Ten serial dilutions (normally 1 µM to 38 pM final
30 concentration) from stock 10 mM solutions of compounds prepared in DMSO. All serial dilutions were made using 20% DMSO. DMSO content in assay is less than 1%. Incubation mixtures were

2004-04-14

- 7 -

prepared in quadruplicate in 96-well plates (Costar). Final assay volumes per well were 10 μ l compound/nonspecific/control (1% DMSO), 20 μ l membranes, 20 μ l [3 H]-Imipramine Hydrochloride, and 150 μ l AB. Specific binding was defined by using 10 μ M Imipramine. The binding reaction was initiated by adding membranes immediately after adding the radioligand to wells containing buffer plus either test compound, nonspecific, or control. The assay plates were placed on a plate shaker and shaken for thirty min while the reactions reached equilibrium. The plates were then filtered through Beckman GF/B filters, presoaked in 6% PEI, using a Packard Filtermate 196. Filters were washed 5x with 0.2 ml ice-cold wash buffer (5 mM Tris HCl, pH 7.4.) The filters were dried and 35 μ l of Microscint20 (Packard) was added to each well. The plates were then counted on a Packard TopCount to determine CPM's per well. Ki values were determined for each test compound utilizing the graphic and analytical software package, GraphPad Prism.

Test B — NK₁ FLIPR Assay using Fluo-4 Dye:

FLIPR assays are performed with a device marketed by Molecular Devices, Inc., designed to precisely measure cellular fluorescence in a high throughput whole-cell assay. (Schroeder et. al., J. Biomolecular Screening, 1(2), p 75-80, 1996).

Compounds were evaluated for potency in blocking the response of U373 cells to the NK₁ receptor agonist Acetyl-[Arg⁶, Sar⁹, Met(O₂)¹¹]-Substance P (ASMSP) using a FLIPR instrument.

U373 cells were loaded with Fluo-4 dye (Molecular Probes) for 45 min at 37 °C and exposed to graded concentrations of compounds for 15 min at room temperature before being challenged with 10 nM – 12 nM ASMSP (an approximately EC₈₀ concentration). Responses were measured as the peak relative fluorescence after agonist addition. pIC₅₀s were calculated from eleven-point concentration-response curves for each compound.

Reagents:

Cell culture medium:

Eagle's MEM with Earle's salts and l-glutamine (500 mL)

Non-essential amino acids, 100 x (5 mL)

Sodium pyruvate, 100 mM (5 mL)

L-Glutamine, 200 mM (5 mL)

FBS (50 mL)

Cellgro 10-010-CV

Cellgro 25-025-CI

Cellgro 25-000-CI

Cellgro 25-005-CI

Cellgro 35-010-CV

Cell harvesting reagents:

101432

2004-04-14

- 8 -

DPBS, 1x without Ca^{++} & Mg^{++}

Cellgro 21-031-CV

1x Trypsin -EDTA (0.5% Trypsin, 0.53% EDTA-4Na)

Cellgro 25-052-CI

Cell plating medium:

UltraCULTURE

BioWhittaker 12-725F

5 L-Glutamine, 200 mM (5 mL/500 mL)

Cellgro 25-005-CI

Working buffer:

10x Hank's balanced salt solution (100 mL/L)

Gibco 14065-056

HEPES buffer 1 M (15 mL/L, [final] 15 mM)

Cellgro 25-060-CI

Probenecid (0.71g dissolved in 6 mL 1 M NaOH for 1L,

10 [final] 2.5 mM)

Sigma P-8761

DDH₂O to 1 L, adjust pH to 7.4 with NaOH

Dye solution:

Fluo-4, AM dye, Molecular Probes F-14201. 50 µg lyophilized dye is dissolved in 23 µL DMSO plus 23 µL Pluronic F-127 (Molecular Probes P-3000). The 46 µL of solubilized fluo-4 dye is

15 then added to 10 mL of working buffer solution to provide a working dye concentration of 5 µM. Each 10 mL of diluted dye is sufficient for a 384-well-plate of cells at 25 µL per well.

Agonist:

Acetyl-[Arg⁶, Sar⁹, Met(O₂)¹¹]-Substance P (ASMSP)

Stock solution of 3.33×10^{-2} M. Dissolve 100 mg in 3.05 mL DMSO and store in aliquots at 4 °C

20

Miscellaneous:

DMSO (to dissolve compounds and for tip wash)

Cell culture and plating procedures:

U373 cells were grown in cell culture medium described above (30 mL per T-150 flask) and harvested when confluent as follows. Medium was removed by aspiration and cells were washed 1x with 12 mL DPBS, without Ca^{++} and Mg^{++} . The DPBS was aspirated and replaced with 3 mL trypsin EDTA. The cells plus trypsin/EDTA were incubated about 2 min at room temperature, until the cells detached from the flask. The harvesting reaction was quenched by addition of 9 mL culture medium and cells were resuspended by trituration.

Cells were passaged at a transfer density of 1:4 every four days. For experiments, cells were counted, pelleted by centrifugation at 400 x g for 5 min and resuspended in cell plating medium at a density of 480,000 cells/mL. 25 µL of this cell suspension was added to each well

of a black-walled 384-well plate (Falcon Microtest, 35 3962) using a Labsystems Multidrop 384 to give 12,000 cells per well. Plates were incubated at 37 °C overnight (minimum 15 h, maximum 23 h) before use.

Compound and agonist preparation:

5 Compounds were dissolved in DMSO at a concentration of 10 mM and 120 µL of these solutions were transferred to the first well (column 1) of each row of a 96-well, round-bottomed, polypropylene storage plate (Costar 3365). Compounds on two such plates were then serially diluted simultaneously in DMSO using a Biomek 2000. 4 µL of each dilution was transferred to a deep well plate (Beckman Coulter 267006) which had been prepared previously to contain 400
10 µL of freshly made working buffer in each well. Concentrations resulting from this procedure are shown in Table 1. The final compound concentrations in the assay span 11 points, between 10 µM and 0.1 nM, in half-log increments.

 The contents of the wells were mixed, and 45 µL of each dilution were transferred - in duplicate - to a 384-well polypropylene compound loading plate (Fisher 12-565-507) so that the
15 384-well plate contained duplicates of each of the compounds from both 96-well plates over the concentration ranges. Columns 23 and 24 of the plate contain no compound and serve as controls. Wells A – N in columns 23 and 24 were loaded with agonist only and therefore represent the maximal response. Wells O – P in columns 23 and 24 were loaded with only buffer, no agonist, and therefore represent the minimum response.

20 An ASMSP agonist loading plate was made by taking stock concentration of ASMSP and diluting in working buffer to give a concentration of 3.3×10^{-8} M. 45 µL of this solution were transferred to all wells of a 384-well polypropylene agonist loading plate (Fisher 12-565-507) except wells O23, O24, P23 & P24 which contained buffer alone and served as unstimulated controls.

25 Dye Loading cells and adding compound:

 For each 384-well assay plate of cells, 10 mL of diluted Fluo-4 dye was prepared as stated above in the methods/reagents section. First, each 384-well cell plate was washed once with working buffer on a CCS Packard plate washer. Any remaining post-wash buffer in the wells was removed by hand and 25 µL per well of Fluo-4 dye was added using a Labsystems Multidrop
30 384. The cell plate was returned to a 37 °C incubator for 45 min to allow the dye to permeate the cells. After 45 min of dye loading, the cell plates were washed twice with working buffer,

leaving a 30 μ L volume of buffer in each well. 5 μ L of compound dilutions were transferred from the compound plate to the cell plate using a PlateMate. Assay plates were incubated in the presence of compound for 15 min at room temperature in the dark, and then loaded onto FLIPR.

Recording responses in FLIPR:

- 5 After the 15 min compound pre-incubation, the plates were loaded onto the FLIPR instrument, 15 μ L of ASMSMSP agonist was added and the cellular response to the agonist was recorded for 90 seconds. The response is measured as the peak relative fluorescence after agonist addition.

Data analysis:

- 10 Results contained in the .stat files generated by FLIPR were pasted into an Excel analysis template and, after outliers were excluded, IC_{50} values were calculated within the template using XLfit. Individual IC_{50} values were reported, along with pIC_{50} . When the two IC_{50} 's obtained for a compound differed by more than 3-fold that compound was assayed one or two more times to re-determine the value.

- 15 Results:

Ki values obtained in the SERT assay for compounds of the invention ranged from less than 2 nM to about 180 nM. IC_{50} values obtained in the FLIPR assay for compounds of the invention ranged from about 70 nM to about 2 μ M.

Examples:

- 20 The invention is illustrated by, but not limited to, the following examples in which descriptions, where applicable and unless otherwise stated, the following terms, abbreviations and conditions are used:

aq., aqueous; atm, atmospheric pressure; BOC, 1,1-dimethylethoxycarbonyl, ACN, acetonitrile; DCM, dichloromethane; DMR, N,N-dimethylformamide; DMSO, dimethyl sulfoxide; EtOH, ethanol; Et₂O, diethyl ether; EtOAc, ethyl acetate; h, hour(s); HPLC, high performance liquid chromatography; HOBT, 1-hydroxybenzotriazole; MeOH, methanol; min, minutes; MS, mass spectrum; NMR, nuclear magnetic resonance; psi, pounds per square inch; RT, room temperature; sat., saturated; TEA, triethylamine; TFA, trifluoroacetic acid; THF, tetrahydrofuran.

- 30 Temperatures are given in degrees Celsius ($^{\circ}$ C); unless otherwise stated, operations were carried out at room or ambient temperature (18-25 $^{\circ}$ C).

Organic solutions were dried over anhydrous sodium or magnesium sulfate; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (4.5-30 mm Hg) with a bath temperature of up to 60 °C.

Chromatography means flash column chromatography on silica gel unless otherwise
5 noted; solvent mixture compositions are given as volume percentages or volume ratios.

When given, NMR data is in the form of delta values for major diagnostic protons (given in parts per million (ppm) relative to tetramethylsilane as an internal standard) determined at 300 MHz.

Melting points are uncorrected.

10 Mass spectra (MS) were obtained using an automated system with atm chemical ionization (APCI) unless otherwise indicated. Masses corresponding to the major isotopic component, or the lowest mass for compounds with multiple masses with nearly equivalent abundance (isotope splitting), are reported.

Where noted that a final compound was converted to the citrate salt, the free base was
15 dissolved in MeOH, DCM, or ACN, combined with citric acid (1.0 equivalents) in MeOH, concentrated under reduced pressure and dried under vacuum (25-60 °C). When indicated that the salt was isolated by filtration from Et₂O, the citrate salt of the compound was stirred in Et₂O for 4-18 h, recovered by filtration, washed with Et₂O, and dried under vacuum (25-60 °C).

Examples:

20 The invention is illustrated by, but not limited to, the following examples in which descriptions, where applicable and unless otherwise stated, the following terms, abbreviations and conditions are used:

aq., aqueous; atm, atmospheric pressure; BOC, 1,1-dimethylethoxycarbonyl, ACN, acetonitrile; DCM, dichloromethane; DMF, N,N-dimethylformamide; DMSO, dimethyl
25 sulfoxide; EtOH, ethanol; Et₂O, diethyl ether; EtOAc, ethyl acetate; h, hour(s); HPLC, high performance liquid chromatography; HOBT, 1-hydroxybenzotriazole; MeOH, methanol; min, minutes; MS, mass spectrum; NMR, nuclear magnetic resonance; psi, pounds per square inch; RT, room temperature; sat., saturated; TEA, triethylamine; TFA, trifluoroacetic acid; THF, tetrahydrofuran.

30 Temperatures are given in degrees Celsius (°C); unless otherwise stated, operations were carried out at room or ambient temperature (18-25 °C).

101432

2004 -04- 14

- 12 -

Organic solutions were dried over anhydrous sodium or magnesium sulfate; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (4.5-30 mm Hg) with a bath temperature of up to 60 °C.

Chromatography means flash column chromatography on silica gel unless otherwise
5 noted; solvent mixture compositions are given as volume percentages or volume ratios.

When given, NMR data is in the form of delta values for major diagnostic protons (given in parts per million (ppm) relative to tetramethylsilane as an internal standard) determined at 300 MHz.

Melting points are uncorrected.

10 Mass spectra (MS) were obtained using an automated system with atm chemical ionization (APCI) unless otherwise indicated. Masses corresponding to the major isotopic component, or the lowest mass for compounds with multiple masses with nearly equivalent abundance (isotope splitting), are reported.

Where noted that a final compound was converted to the citrate salt, the free base was
15 dissolved in MeOH, DCM, or ACN, combined with citric acid (1.0 equivalents) in MeOH, concentrated under reduced pressure and dried under vacuum (25-60 °C). When indicated that the salt was isolated by filtration from Et₂O, the citrate salt of the compound was stirred in Et₂O for 4-18 h, recovered by filtration, washed with Et₂O, and dried under vacuum (25-60 °C).

Example 1: N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-2-(4-fluorophenyl)-N-methyl-2-pyrrolidin-1-ylacetamide.
20

(4-Fluorophenyl)(pyrrolidin-1-yl)acetic acid (400 mg, 1.79 mmol), 3-methoxy-4-[(methylamino)methyl]-2-naphthonitrile (405 mg, 1.79 mmol), HOBt (363 mg, 2.69 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (412 mg, 2.15 mmol) were reacted together in DCM (15 mL) at RT overnight. The reaction mixture was partitioned between
25 water (20 mL) and DCM (20 mL). The organic layer was washed with sat. aq. NaHCO₃, dried over Na₂SO₄, filtered through a pad of diatomaceous earth and the volatiles were removed under reduced pressure. Chromatography of the residue on SiO₂ (35 g) (0-5% MeOH:DCM) afforded the title compound as a white solid (300 mg). ¹H NMR (300.1 MHz, DMSO) δ 8.61 (s, 6H), 8.04 (d, J = 8.5 Hz, 7H), 7.91 (d, J = 8.5 Hz, 7H), 7.71 - 7.58 (m, 26H), 7.29 (t, J = 8.5 Hz, 12H), 5.70 (s, 6H), 5.24 (d, J = 14.3 Hz, 6H), 5.00 (d, J = 14.3 Hz, 6H), 3.98 (s, 18H), 3.78 - 3.72 (m, 7H),
30 3.23 - 3.14 (m, 6H), 2.99 - 2.90 (m, 7H), 2.82 - 2.76 (m, 7H), 2.61 (s, 19H), 2.06 - 1.96 (m, 19H),

1.90 - 1.78 (m, 8H). The citrate salt was formed by the addition of citric acid (1.0 equivalents) to a methanolic solution of the title compound. Concentration under reduced pressure afforded the desired salt form of the product as a white foam. MS m/z 432.2 (M+H)⁺.

The requisite (4-fluorophenyl)(pyrrolidin-1-yl)acetic acid was synthesized using the following method.

Pyrrolidine (200 mg, 2.8 mmol), glyoxylic acid monohydrate (258 mg, 2.8 mmol) and (4-fluorophenyl)boronic acid (393 mg, 2.8 mmol) were reacted together in DCM (10 mL) at reflux overnight. The volatiles were removed under reduced pressure, the resulting residue was dissolved in DCM (1 mL). The solution was added to vigorously stirred Et₂O (100 mL) to afford the product as a pale orange powder (430 mg). MS m/z 224.1 (M+H)⁺.

The requisite 3-methoxy-4-[(methylamino)methyl]-2-naphthonitrile was synthesized using the following method.

Methylamine (500 mL of a 2.0 M solution in MeOH) was added to 4-(iodomethyl)-3-methoxy-2-naphthonitrile (5.00 g, 15.5 mmol). After reacting at RT overnight in a sealed vessel, the volatiles were removed under reduced pressure. The residue was taken up in DCM (350 mL) and washed with sat. aq. NaHCO₃ (550 mL). The organic layer was dried over Na₂SO₄, filtered through a pad of diatomaceous earth and the volatiles were removed under reduced pressure to afford the title compound as a pale yellow powder (3.50 g, 100%). ¹H NMR (300.1 MHz, DMSO) δ 8.53 (s, 1H), 8.26 (d, J = 8.4 Hz, 1H), 8.01 (d, J = 8.4 Hz, 1H), 7.72 (ddd, J = 8.4, 7.0, 1.4 Hz, 1H), 7.62 - 7.56 (m, 1H), 4.06 (s, 2H), 3.94 (s, 3H), 3.31 (s, 1H), 2.38 (s, 3H).

Examples 2-24:

The following compounds were prepared in a manner analogous to that described in Example 1:

Example No.	Name	MW	MS m/z (M+H) ⁺
2	N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-2-(4-fluorophenyl)-N-methyl-2-[3-(methylamino)pyrrolidin-1-yl]acetamide	460.56	461.3
3	N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-2-(4-fluorophenyl)-N-methyl-2-[methyl(pyrrolidin-3-yl)amino]acetamide	460.56	461.3


4	N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-2-(4-fluorophenyl)-N-methyl-2-[(1-methylpyrrolidin-3-yl)amino]acetamide	460.56	461.3
5	N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-2-[3-(dimethylamino)azetidin-1-yl]-2-(4-fluorophenyl)-N-methylacetamide	460.56	461.1
6	2-[azetidin-3-yl(methyl)amino]-N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-2-(4-fluorophenyl)-N-methylacetamide	446.53	447.3
7	2-(3-aminopyrrolidin-1-yl)-N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-2-(4-fluorophenyl)-N-methylacetamide	446.53	447.1
8	2-(3-aminoazetidin-1-yl)-N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-2-(4-fluorophenyl)-N-methylacetamide	432.50	433.2
9	N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-2-(3,4-difluorophenyl)-N-methyl-2-pyrrolidin-1-ylacetamide	449.5	450.0
10	N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-2-(3-methoxyphenyl)-N-methyl-2-pyrrolidin-1-ylacetamide	443.54	444.1
11	N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-N-methyl-2-phenyl-2-pyrrolidin-1-ylacetamide	413.52	414.1
12	N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-2-(3-fluorophenyl)-N-methyl-2-pyrrolidin-1-ylacetamide	431.51	432.1
13	2-(3-chloro-4-fluorophenyl)-N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-N-methyl-2-pyrrolidin-1-ylacetamide	465.95	466.1
14	2-(3,4-difluorophenyl)-N-[2-methoxy-5-(trifluoromethoxy)benzyl]-N-methyl-2-pyrrolidin-1-ylacetamide	458.43	459.1
15	2-(3-chloro-4-fluorophenyl)-N-[2-methoxy-5-(trifluoromethoxy)benzyl]-N-methyl-2-pyrrolidin-1-ylacetamide	474.88	475.0
16	2-(4-fluorophenyl)-N-[2-methoxy-5-(trifluoromethoxy)benzyl]-N-methyl-2-pyrrolidin-1-ylacetamide	440.44	441.1
17	N-{2-[3,5-bis(trifluoromethyl)phenyl]ethyl}-2-(3-methoxyphenyl)-N-methyl-2-pyrrolidin-1-ylacetamide	488.47	489.1
18	N-{2-[3,5-bis(trifluoromethyl)phenyl]ethyl}-N-methyl-2-phenyl-2-pyrrolidin-1-ylacetamide	458.44	459.4
19	N-{2-[3,5-bis(trifluoromethyl)phenyl]ethyl}-2-(4-fluorophenyl)-N-methyl-2-pyrrolidin-1-ylacetamide	476.43	477.4
20	2-azetidin-1-yl-N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-2-(4-fluorophenyl)-N-methylacetamide	417.48	418.3
21	N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-2-(4-fluorophenyl)-N-methyl-2-piperidin-1-ylacetamide	445.54	446.4

22	N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-2-(dimethylamino)-2-(4-fluorophenyl)-N-methylacetamide	405.47	406.4
23	N-[(3-cyano-2-methoxy-1-naphthyl)methyl]-2-(4-fluorophenyl)-N-methyl-2-morpholin-4-ylacetamide	447.51	448.1
24	4-{[2-(4-fluorophenyl)-2-pyrrolidin-1-ylethoxy]methyl}-3-methoxy-2-naphthonitrile	404.48	405.4



Claims

1. A compound according to structural diagram I:



I

wherein:

R^1 and R^2 are independently selected from C_{1-6} alkyl or C_{1-6} alkenyl, or together with the N to which they are bound, form heterocycle containing 3-6 atoms or heterocycle substituted with moieties independently selected from hydrogen, halogen, C_{1-4} alkyl, C_{1-4} alkoxy or C_{1-4} alkyl substituted with 1, 2 or 3 halo moieties, amino, or amino substituted with C_{1-4} alkyl, C_{1-4} alkoxy or C_{1-4} alkyl, substituted with 0, 1, 2 or 3 halo moieties; in vivo-hydrolysable precursors thereof, and pharmaceutically-acceptable salts thereof;

R^3 is independently selected from hydrogen or C_{1-6} alkyl;

X is selected from O or NR^4 ;

R^4 is selected from hydrogen or C_{1-6} alkyl;

Y is C_{1-6} alkyl or $C(O)$;

n is 0, 1 or 2;

Ar^1 is phenyl or phenyl substituted with moieties independently selected from hydrogen, halogen, C_{1-4} alkyl, C_{1-4} alkoxy or C_{1-4} alkyl substituted with 1, 2 or 3 halo moieties; in vivo-hydrolysable precursors thereof, and pharmaceutically-acceptable salts thereof; and

Ar^2 phenyl, naphthyl, tetralin, or phenyl, naphthyl or tetralin substituted with moieties independently selected from hydrogen, halogen, cyano, nitro, C_{1-4} alkyl, C_{1-4} alkoxy or C_{1-4} alkyl substituted with 1, 2 or 3 halo moieties; in vivo-hydrolysable precursors thereof, and pharmaceutically-acceptable salts thereof.

2. Pharmaceutically-acceptable salts of a compound according to Claim 1 made with an inorganic or organic acid which affords a physiologically-acceptable anion.


3. Pharmaceutically-acceptable salts according to Claim 2, wherein said inorganic or organic acid is selected from hydrochloric, hydrobromic, sulfuric, phosphoric, methanesulfonic, sulfamic, para-toluenesulfonic, acetic, citric, lactic, tartaric, malonic, fumaric, ethanesulfonic, benzenesulfonic, cyclohexylsulfamic, salicylic and quinic acids.
- 5 4. A pharmaceutical composition comprising a compound according to Claim 1, an in vivo-hydrolysable precursor or a pharmaceutically-acceptable salt thereof and a pharmaceutically-acceptable carrier.
- 10 5. A method of treating a disease condition wherein antagonism of NK_1 receptors in combination with SRI activity is beneficial which method comprises administering to a warm-blooded animal an effective amount of a compound according to Claim 1 or an in vivo-hydrolysable precursor or a pharmaceutically-acceptable salt thereof.
- 15 6. A method of treating a disease condition wherein antagonism of NK_1 receptors is beneficial which method comprises administering to a warm-blooded animal an effective amount of a compound according to Claim 1 or an in vivo-hydrolysable precursor or a pharmaceutically-acceptable salt thereof.
- 20 7. A method of treating a disease condition wherein SRI activity is beneficial with method comprises administering to a warm-blooded animal an effective amount of a compound according to Claim 1 or a in vivo-hydrolysable precursor or a pharmaceutically-acceptable salt thereof.
- 25 8. The use of a compound according to Claim 1 or an in vivo-hydrolysable precursor or a pharmaceutically-acceptable salt thereof in the preparation of a medicament for use in a disease condition wherein antagonism of the NK_1 receptors or SRI activity is beneficial.
- 30 9. The use of a compound according to Claim 1 or an in vivo-hydrolysable precursor or a pharmaceutically-acceptable salt thereof in the preparation of a medicament for use in a disease condition wherein antagonism of the NK_1 receptors and SRI activity is beneficial.

10. A method for treating a disorder or condition selected from hypertension, depression in cancer patients, depression in Parkinson's patients, postmyocardial infarction depression, subsyndromal symptomatic depression, depression in infertile women, pediatric depression, major depression, single episode depression, recurrent depression, child abuse induced depression, post partum depression, generalized anxiety disorder, agoraphobia, social phobia, simple phobias, posttraumatic stress syndrome, avoidant personality disorder, premature ejaculation, anorexia nervosa, bulimia nervosa, obesity, addictions to alcohol, cocaine, heroin, phenobarbital, nicotine or benzodiazepines; cluster headache, migraine, pain, Alzheimer's disease, obsessive-compulsive disorder, panic disorder, dementia, amnesic disorders, age-related cognitive decline, dementia in Parkinson's disease, neuroleptic-induced parkinsonism, tardive dyskinesias, hyperprolactinaemia, vasospasm, cerebral vasculature vasospasm, cerebellar ataxia, gastrointestinal tract disorders, negative symptoms of schizophrenia, premenstrual syndrome, fibromyalgia syndrome, stress incontinence, Tourette's syndrome, trichotillomania, kleptomania, male impotence, attention deficit hyperactivity disorder, chronic paroxysmal hemicrania and headache associated with vascular disorders in a mammal, wherein antagonism of the NK₁ receptors and SRI activity is beneficial, comprising administering an effective amount of a compound according to Claim 1 or a pharmaceutically-acceptable salt thereof effective in treating such disorder or condition.
11. The method according to any one of Claims 5, 6 or 7, wherein said compound is administered in combination with a pharmaceutically-acceptable carrier.

A B S T R A C T

Title: ARYL GLYCINE DERIVATIVES AND THEIR USE

- 5 Compounds of the following formula



I

- wherein R^1 , R^2 , R^3 , n , X , Y , Ar^1 and Ar^2 are as defined in the specification, in vivo-hydrolysable precursors thereof, pharmaceutically-acceptable salts thereof, the use in therapy and
- 10 pharmaceutical compositions and methods of treatment using the same.